




# 2.  $\rightarrow q_0 \xrightarrow{a/a/R} q_1$   $\delta(q_0, a) = (q_1, a, R)$   
 $Q = \{q_0, q_1\}$ ,  $F = \{q_1\}$ ,  $\Sigma = \{a, b\}$ ,  $\Gamma = \{a, b, \square\}$

# 3.  $aba$ :  
 $q_0 aba \vdash x q_1 ba \vdash q_2 x ya \vdash x q_0 ya \vdash x y q_3 a$


# 4. No.

# 5.  $L(\underline{a}b^* + \underline{b}\underline{b}^*\underline{a}(\underline{a} + \underline{b})^*)$


# 6. The Turing Machine halts in a non-accepting state



#9.



#11(a) Modify the machine in Example 9.10 to write two  $i$ 's for each  $x$  :



Section 11.1 #12 p. 282.

- 12 (a) If  $L_1$  is recursive and  $L_2$  is r.e., then  $L_2 - L_1$  is necessarily r.e. because  $\bar{L}$  is r.e.,  
 So  $L_2 - L_1 = L_2 \cap \bar{L}_1 =$  intersection of two r.e. sets = r.e.
- (b) If  $L_1$  is recursive and  $L_2$  is r.e., then  $L_1 - L_2$  is not necessarily r.e. because if we take  $L_1 = \Sigma^*$  and  $L_2 =$  a non-recursive r.e. subset of  $\Sigma^*$ , then  $L_1 - L_2 =$  a non-r.e. subset of  $\Sigma^*$ .

- #2 The set of all r.e. languages is countable since every r.e. language is associated with a TM and the number of different TM's is countable. If there were countably many non-r.e. languages as well, then there would be countably many languages altogether. But the set of all languages on  $\Sigma$  is uncountable because  $\mathcal{L}(\Sigma) = \mathcal{P}(\Sigma^*) \approx \mathcal{P}(\mathbb{N})$  which is uncountable. So there are uncountably many non-r.e. languages.
- #5. Suppose  $\bar{L}$  is recursive. Then  $\bar{\bar{L}}$  will also be recursive by the closure theorem for recursive languages. But  $\bar{\bar{L}} = L$ . So  $L$  is recursive. Hence  $L$  is r.e. But we were told that  $L$  was not r.e. So if  $L$  is not r.e., then  $\bar{L}$  cannot be recursive.
- #6 Suppose  $L_1$  and  $L_2$  are r.e. Let  $M_1$  and  $M_2$  be TM's such that  $L(M_1) = L_1$  and  $L(M_2) = L_2$ . A third TM  $M$  can run both  $M_1$  and  $M_2$  together and enter and enter a final state if either one does (e.g.,  $M$  can put  $M_1$  &  $M_2$  through their respective moves alternately). Any  $w$  which causes either  $M_1$  or  $M_2$  to enter a final state will cause  $M$  to enter a final state and vice versa.  $\therefore L(M_1) \cup L(M_2) = L(M)$   
 $\therefore L_1 \cup L_2$  will be r.e.

#7 Yes. Let  $M_1$  and  $M_2$  be as in problem 5 above. Another TM  $M$  can be designed to run both together and enter a final state if both  $M_1$  and  $M_2$  do. (For example,  $M$  can simulate  $M_1$  and  $M_2$  as follows. Put  $M_1$  &  $M_2$  alternately through their moves and if one of them halts in a final state, then continue only with the other one until it halts in a final state.) Then  $L(M) = L(M_1) \cap L(M_2)$ . So  $L_1 \cap L_2$  is also r.e.

#8 If  $L_1$  &  $L_2$  are recursive, then  $\bar{L}_1, \bar{L}_2, L_1$  and  $L_2$  will all be r.e. So by problems 5 and 6,

$L_1 \cup L_2$  will be r.e.  
and  $\bar{L}_1 \cap \bar{L}_2 = \overline{L_1 \cup L_2}$  will be r.e.  
Since  $L_1 \cup L_2$  &  $\overline{L_1 \cup L_2}$  are both r.e., it follows that  $L_1 \cup L_2$  is recursive.

Similarly  $L_1 \cap L_2$  &  $\overline{L_1 \cap L_2} = \bar{L}_1 \cup \bar{L}_2$  will be both r.e. Hence  $L_1 \cap L_2$  will be recursive.

Note: recursive sets are closed under  $\cup, \cap$  and compliments. r.e. sets are not closed under compliments.

$$\begin{aligned}
 \#1. (a) \text{ ADD}(3,4) &= \text{ADD}(3,3+1) = \text{ADD}(3,3) + 1 \\
 &= \text{ADD}(3,2+1) + 1 = (\text{ADD}(3,2) + 1) + 1 \\
 &= (\text{ADD}(3,1+1) + 1) + 1 = ((\text{ADD}(3,1) + 1) + 1) + 1 \\
 &= ((\text{ADD}(3,0+1) + 1) + 1) + 1 \\
 &= (((\text{ADD}(3,0) + 1) + 1) + 1) + 1 \\
 &= (((3+1) + 1) + 1) + 1 = ((4+1) + 1) + 1 \\
 &= (5+1) + 1 = 6+1 = 7
 \end{aligned}$$

$$\begin{aligned}
 (b) \text{ MULT}(2,3) &= \text{MULT}(2,2+1) = \text{MULT}(2,2) + 2 \\
 &= \text{MULT}(2,1+1) + 2 = (\text{MULT}(2,1) + 2) + 2 \\
 &= (\text{MULT}(2,0+1) + 2) + 2 = ((\text{MULT}(2,0) + 2) + 2) + 2 \\
 &= ((0+2) + 2) + 2 = \dots = 6
 \end{aligned}$$

$$\#2 \text{ Note } x \dot{-} ((x \dot{-} y) + (y \dot{-} x)) = \begin{cases} 1 & \text{if } x=y \\ 0 & \text{if } x \neq y \end{cases}$$

$\therefore$  equal  $(x, y)$

$$= \text{monus}(c_1, (p_1(x, y), \text{add}(\text{monus}(x, y), \text{monus}(p_2(x, y), p_1(x, y))))))$$

~~#2~~ (a) Note that  $f(x, y) = x \dot{-} h(x, y)$  where

$$h(x, y) = \begin{cases} 0 & \text{if } x \neq y \\ x & \text{if } x = y \end{cases}$$

$$\begin{aligned}
 \text{So } f(x, y) &= x \dot{-} (x \cdot \text{equal}(x, y)) \\
 &= \text{monus}(p_1(x, y), \text{mult}(p_1(x, y), \text{equal}(x, y)))
 \end{aligned}$$

$$(b) f(0) = 1$$

$$f(y+1) = \text{mult}(y, f(y))$$

From this we can see  $f(y) = y!$  is primitive recursive by doing a little work.