

What happens when the exponents of the D.E. are imaginary?

$x^2 y'' + xy' + y = 0$ (This can be solved as a Cauchy-Euler equation, but let's use the method of Frobenius) First, note that 0 is a regular singular point.

Assume a solution of the form $y = \sum_{n=0}^{\infty} C_n x^{n+r}$, $C_0 \neq 0$, $x > 0$.

Substituting $y, y',$ and y'' in the D.E. yields:

$$\sum_{n=0}^{\infty} (n+r-1)(n+r)C_n x^{n+r} + \sum_{n=0}^{\infty} (n+r)C_n x^{n+r} + \sum_{n=0}^{\infty} C_n x^{n+r} = 0$$

Setting $n=0$ to get the indicial equation $\rightarrow (r-1)r+r+1=0$

$$r^2 + 1 = 0$$

$$r = \pm i$$

For $n \geq 1$ $(n+r-1)(n+r)C_n + (n+r)C_n + C_n = 0$

$$(n+r)[n+r-1+1]C_n + C_n = 0$$

$$[(n+r)^2 + 1]C_n = 0$$

Case I: $r = i$

$$[(n+i)^2 + 1]C_n = 0$$

$$[n^2 + 2in]C_n = 0$$

$$n(n+2i)C_n = 0$$

Since $n \geq 1$, $n \neq 0$ and $n \neq -2i$. So $C_n = 0$

Thus all C_i except C_0 are zero.

We get the solution $y_1 = C_0 x^i$

Case II: $r = -i$

$$[(n-i)^2 + 1]C_n = 0$$

$$[n^2 - 2in]C_n = 0$$

$$n(n-2i)C_n = 0$$

Since $n \geq 1$, $n \neq 0$ and $n \neq 2i$. So $C_n = 0$

Thus all C_i except C_0 are zero.

We get the solution $y_2 = C_0^* x^{-i}$

Using the identity $e^{ix} = \cos x + i \sin x$:

$$x^i = (e^{\ln x})^i = e^{i \ln x} = \cos(\ln x) + i \sin(\ln x)$$

$$x^{-i} = (e^{\ln x})^{-i} = e^{-i \ln x} = \cos(-\ln x) + i \sin(-\ln x) = \cos(\ln x) - i \sin(\ln x)$$

$x^i + x^{-i} = 2 \cos(\ln x) \Rightarrow \cos(\ln x) = \frac{1}{2} x^i + \frac{1}{2} x^{-i} \Rightarrow \cos(\ln x)$ is a linear combination of solutions $\Rightarrow \cos(\ln x)$ is a solution.

$x^i - x^{-i} = 2i \sin(\ln x) \Rightarrow \sin(\ln x) = \frac{1}{2i} x^i - \frac{1}{2i} x^{-i} \Rightarrow \sin(\ln x)$ is a linear combination of solutions $\Rightarrow \sin(\ln x)$ is a solution.

Since $\cos(\ln x)$ and $\sin(\ln x)$ are linearly independent (verify using Wronskian) on $(0, +\infty)$, the general solution can be written $y = C_1 \cos(\ln x) + C_2 \sin(\ln x)$