

Problem 1.7

From Eq. 1.33, $\frac{d\langle p \rangle}{dt} = -i\hbar \int \frac{\partial}{\partial t} (\Psi^* \frac{\partial \Psi}{\partial x}) dx$. But, noting that $\frac{\partial^2 \Psi}{\partial x \partial t} = \frac{\partial^2 \Psi}{\partial t \partial x}$ and using Eqs. 1.23-1.24:

$$\begin{aligned} \frac{\partial}{\partial t} \left(\Psi^* \frac{\partial \Psi}{\partial x} \right) &= \frac{\partial \Psi^*}{\partial t} \frac{\partial \Psi}{\partial x} + \Psi^* \frac{\partial}{\partial x} \left(\frac{\partial \Psi}{\partial t} \right) = \left[-\frac{i\hbar}{2m} \frac{\partial^2 \Psi^*}{\partial x^2} + \frac{i}{\hbar} V \Psi^* \right] \frac{\partial \Psi}{\partial x} + \Psi^* \frac{\partial}{\partial x} \left[\frac{i\hbar}{2m} \frac{\partial^2 \Psi}{\partial x^2} - \frac{i}{\hbar} V \Psi \right] \\ &= \frac{i\hbar}{2m} \left[\Psi^* \frac{\partial^3 \Psi}{\partial x^3} - \frac{\partial^2 \Psi^*}{\partial x^2} \frac{\partial \Psi}{\partial x} \right] + \frac{i}{\hbar} \left[V \Psi^* \frac{\partial \Psi}{\partial x} - \Psi^* \frac{\partial}{\partial x} (V \Psi) \right] \end{aligned}$$

The first term integrates to zero, using integration by parts twice, and the second term can be simplified to $V \Psi^* \frac{\partial \Psi}{\partial x} - \Psi^* V \frac{\partial \Psi}{\partial x} - \Psi^* \frac{\partial V}{\partial x} \Psi = -|\Psi|^2 \frac{\partial V}{\partial x}$. So

$$\frac{d\langle p \rangle}{dt} = -i\hbar \left(\frac{i}{\hbar} \right) \int -|\Psi|^2 \frac{\partial V}{\partial x} dx = \left\langle -\frac{\partial V}{\partial x} \right\rangle. \quad \text{QED}$$

Problem 1.9

(a)

$$1 = 2|A|^2 \int_0^\infty e^{-2amx^2/\hbar} dx = 2|A|^2 \frac{1}{2} \sqrt{\frac{\pi}{2am/\hbar}} = |A|^2 \sqrt{\frac{\pi\hbar}{2am}}; \quad \boxed{A = \left(\frac{2am}{\pi\hbar}\right)^{1/4}}$$

(b)

$$\frac{\partial\Psi}{\partial t} = -ia\Psi; \quad \frac{\partial\Psi}{\partial x} = -\frac{2amx}{\hbar}\Psi; \quad \frac{\partial^2\Psi}{\partial x^2} = -\frac{2am}{\hbar} \left(\Psi + x\frac{\partial\Psi}{\partial x}\right) = -\frac{2am}{\hbar} \left(1 - \frac{2amx^2}{\hbar}\right)\Psi.$$

Plug these into the Schrödinger equation, $i\hbar\frac{\partial\Psi}{\partial t} = -\frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial x^2} + V\Psi$:

$$\begin{aligned} V\Psi &= i\hbar(-ia)\Psi + \frac{\hbar^2}{2m} \left(-\frac{2am}{\hbar}\right) \left(1 - \frac{2amx^2}{\hbar}\right)\Psi \\ &= \left[\hbar a - \hbar a \left(1 - \frac{2amx^2}{\hbar}\right) \right] \Psi = 2a^2mx^2\Psi, \quad \text{so } \boxed{V(x) = 2ma^2x^2}. \end{aligned}$$

(c)

$$\langle x \rangle = \int_{-\infty}^{\infty} x|\Psi|^2 dx = \boxed{0}. \quad \text{[Odd integrand.]}$$

$$\langle x^2 \rangle = 2|A|^2 \int_0^\infty x^2 e^{-2amx^2/\hbar} dx = 2|A|^2 \frac{1}{2^2(2am/\hbar)} \sqrt{\frac{\pi\hbar}{2am}} = \boxed{\frac{\hbar}{4am}}.$$

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = \boxed{0}.$$

$$\begin{aligned} \langle p^2 \rangle &= \int \Psi^* \left(\frac{\hbar}{i} \frac{\partial}{\partial x}\right)^2 \Psi dx = -\hbar^2 \int \Psi^* \frac{\partial^2\Psi}{\partial x^2} dx \\ &= -\hbar^2 \int \Psi^* \left[-\frac{2am}{\hbar} \left(1 - \frac{2amx^2}{\hbar}\right)\Psi\right] dx = 2am\hbar \left\{ \int |\Psi|^2 dx - \frac{2am}{\hbar} \int x^2 |\Psi|^2 dx \right\} \\ &= 2am\hbar \left(1 - \frac{2am}{\hbar} \langle x^2 \rangle\right) = 2am\hbar \left(1 - \frac{2am}{\hbar} \frac{\hbar}{4am}\right) = 2am\hbar \left(\frac{1}{2}\right) = \boxed{am\hbar}. \end{aligned}$$

(d)

$$\sigma_x^2 = \langle x^2 \rangle - \langle x \rangle^2 = \frac{\hbar}{4am} \Rightarrow \boxed{\sigma_x = \sqrt{\frac{\hbar}{4am}}}; \quad \sigma_p^2 = \langle p^2 \rangle - \langle p \rangle^2 = am\hbar \Rightarrow \boxed{\sigma_p = \sqrt{am\hbar}}.$$

$\sigma_x \sigma_p = \sqrt{\frac{\hbar}{4am}} \sqrt{am\hbar} = \frac{\hbar}{2}$. This is (just barely) consistent with the uncertainty principle.

Problem 1.15

(a) Eq. 1.24 now reads $\frac{\partial \Psi^*}{\partial t} = -\frac{i\hbar}{2m} \frac{\partial^2 \Psi^*}{\partial x^2} + \frac{i}{\hbar} V^* \Psi^*$, and Eq. 1.25 picks up an extra term:

$$\frac{\partial}{\partial t} |\Psi|^2 = \dots + \frac{i}{\hbar} |\Psi|^2 (V^* - V) = \dots + \frac{i}{\hbar} |\Psi|^2 (V_0 + i\Gamma - V_0 + i\Gamma) = \dots - \frac{2\Gamma}{\hbar} |\Psi|^2,$$

and Eq. 1.27 becomes $\frac{dP}{dt} = -\frac{2\Gamma}{\hbar} \int_{-\infty}^{\infty} |\Psi|^2 dx = -\frac{2\Gamma}{\hbar} P$. QED

(b)

$$\frac{dP}{P} = -\frac{2\Gamma}{\hbar} dt \implies \ln P = -\frac{2\Gamma}{\hbar} t + \text{constant} \implies \boxed{P(t) = P(0)e^{-2\Gamma t/\hbar}}, \text{ so } \boxed{\tau = \frac{\hbar}{2\Gamma}}.$$